



Fig. 1.





■ α_51 casein sequence, promoter or 3' untranslated region.

2 3 The boxes represent the exons in the α -glucosidase sequence, the thin line represents the intron sequences.


The numbers above the boxes are the exon numbers

pA = polyadenylation signal.

ATG = translation initiation site,

TAG = translation stop codon

Fig. 3A.



201050 "TISHOOT"

Fig. 3B.





Fig. 3.C.



60F 29




Fig. 4. A.

Fig. 4. B.






Fig. 5.



90F 29

Fig. 6.



10 OF 29

Fig. 7.




Fig. 8.




Fig. 9.




Fig. 10.




Fig. 11. A.




Fig. 11. B.




Fig. 11. C.




Fig. 11. D.




Fig. 12.




Fig. 13. A.

transgenic whey




Fig. 13. B.

non-transgenic whey






Fig. 14.



_____ 12099801:1_UV1_280nm
 ----- 12099801:1_pH
 -x-x-x- 12099801:1_Cond%
 12099801:1_Fractions

Fig. 15.




12099802:11_UV1_280nm

12099802:11_pH

12099802:11_Cond%

12099802:11_Fractions


Fig. 16.



12099803:12_Fractions

20 of 29

Fig. 17.




12099804:13_UV1_280nm

12099804:13_pH

12099804:13_Cond%

12099804:13_Fractions

Fig. 18.



121099805:1_UV1_280nm


121099805:1_pH

121099805:1_Conc%

121099805:1_Fractions

22 of 29

Fig. 19.




121099806:1_UV1_280nm

121099806:1_pH

121099806:1_Cond%

121099806:1_Fractions

Fig. 20.



—— hatypei01:1_UV1_280nm

----- hatypei01:1_pH

-x-x-x- hatypei01:1_Conc%

hatypei01:1_Fractions

24 of 29

Fig. 21.






Fig. 22.



— hatypei03:12_UV1_280nm
 - - - hatypei03:12_pH
 x x x hatypei03:12_Cond%

hatypei03:12_Fractions

Fig. 23.



— hatypei04:13_UV1_280nm

----- hatypei04:13_pH

-x-x-x- hatypei04:13_Conc%

hatypei04:13_Fractions

Fig. 24.





Fig. 25.



28 OF 29

Fig. 26.

XK16/15 80°C
 cHT type I 10mM Napi pH 6.5 ; QFF eluate
 Run 02249901/02259901/02269901

